



Early View

Editorial

Managing the supportive care needs of those affected by COVID-19

S. Bajwah, A. Wilcock, R. Towers, M. Costantini, C. Bausewein, S. T. Simon, E. Bendstrup, W. Prentice, M. J. Johnson, D. C. Currow, M. Kreuter, A. U. Wells, S. S. Birring, P. Edmonds, I. J. Higginson

Please cite this article as: Bajwah S, Wilcock A, Towers R, *et al*. Managing the supportive care needs of those affected by COVID-19. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.00815-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

Managing the supportive care needs of those affected by COVID-19

Bajwah S, Wilcock A, Towers R, Costantini M, Bausewein C, Simon ST, Bendstrup E, Prentice W, Johnson MJ, Currow DC, Kreuter M, Wells AU, Birring SS, Edmonds P, Higginson IJ

Corresponding Author:

Dr Sabrina Bajwah, Clinical Senior Lecturer Palliative Care, Cicely Saunders Institute, Department of Palliative Care Policy and Rehabilitation, King's College London, Bessemer Road, London SE5 9PJ, UK.

Correspondence to: Email: sabrina.bajwah@kcl.ac.uk Telephone: 020 7848 5826

Dr Andrew Wilcock, Reader in Medical Oncology and Palliative Medicine, University of Nottingham, Nottingham, UK

Dr Richard Towers, Matron Palliative Care, Department of Palliative Care, King's College Hospital, London, UK

Dr Massimo Costantini, Scientific Director, Azienda USL-IRCCS di Reggio Emilia, Italy

Professor Claudia Bausewein, Professor of Palliative Medicine, Department of Palliative Medicine, LMU Klinikum, Ludwig-Maximilians-University, Germany

Dr Steffen T Simon, University of Cologne, Faculty of Medicine and University Hospital, Department of Palliative Medicine and Center for Integrated Oncology, Germany

Dr Elisabeth Bendstrup, Centre for Rare Lung Diseases, Department of Respiratory Diseases and Allergy, Aarhus University Hospital, Denmark

Dr Wendy Prentice, Consultant Palliative Care, Department of Palliative Care, King's College Hospital, London, UK

Professor Miriam J Johnson, Professor of Palliative Care, Wolfson Palliative Care Research Centre, Hull York Medical School, University of Hull, Hull, UK.

Professor David C Currow, Professor of Palliative Care, IMPACCT, Faculty of Health, University of Technology Sydney, Ultimo. Australia,

Professor Michael Kreuter, Center for interstitial and rare lung diseases, Thoraxklinik, University of Heidelberg, Germany and German Center for Lung Research, Heidelberg, Germany

Professor Athol U Wells, Professor of Interstitial Lung Disease, Royal Brompton Hospital, London, UK

Professor Surinder S Birring, Professor of Respiratory Medicine, Centre for Human & Applied Physiological Sciences, School of Basic & Medical Biosciences, Faculty of Life Sciences & Medicine, King's College London, London

Dr Polly Edmonds, Consultant Palliative Care, Department of Palliative Care, King's College Hospital, London, UK

Professor Irene J Higginson, Professor of Palliative Care, Cicely Saunders Institute,
Department of Palliative Care Policy and Rehabilitation, King's College London, UK

Globally, the number of people affected by coronavirus disease 2019 (COVID-19) is rapidly increasing. In most (>80%), the illness is relatively mild and can be self-managed out of hospital. However, in about 20% the illness causes respiratory compromise severe enough to require hospital admission.[1] Patients with severe and critical disease need full active treatment. This may include oxygen for hypoxaemia and ventilatory support, along with optimal management of complications, e.g. super-imposed bacterial infection, and any underlying co-morbidities, e.g. chronic obstructive pulmonary disease, congestive heart failure. To date, no antiviral agent has shown to be effective in treating the disease.[2]

Patients with severe disease not considered suitable for escalation to intensive care, i.e. those who are frail or have multiple co-morbidities, are at very high risk of dying, with an estimated death rate of 15–22%.[3, 4] We have a moral obligation to provide good symptom control to prevent avoidable suffering. Thus, comprehensive care of the patient with COVID-19 requires identification of patients at increased risk of dying, who would benefit from a parallel approach to management. This encompasses optimal symptom management for those with severe disease but who will survive, and expert symptom management and end of life care for those that are deteriorating and in their last days–hours of life. The aim of this editorial is to provide a succinct informative overview to guide respiratory healthcare professionals on the frontline.

Symptom relief

The most common symptoms are breathlessness, cough and fever.

Breathlessness (5–65%)[5-9]: The highest incidence will be in those with severe disease and is expected in those actively dying. The primary driver of breathlessness is the viral lung infection causing an interstitial pneumonia with a reduction in lung diffusing capacity; in some patients this evolves to Acute Respiratory Distress Syndrome (ARDS). The experience of breathlessness is also influenced by emotional, environmental, cultural and social factors,

and optimal management requires a holistic approach. These include non-pharmacological and pharmacological approaches. Non-pharmacological approaches include breathing techniques.[10] Although an electric hand-held fan directed at the face is helpful in other settings [10, 11], this is not recommended in COVID-19 because of the theoretical infection control risk of spreading infected droplets. As an alternative, use of facial cooling with wet wipes (binning after each use, as for tissues) can be tried.

Pharmacological approaches are the mainstay of management for patients with severe disease who are likely to have rapidly worsening breathlessness at rest. Morphine is the opioid of choice.[12, 13] However, alternative strong opioids can be used. Short-acting oral opioids, given as required, may suffice when breathlessness is mild. However, in severe disease or at the end of life, continuous infusions of parenteral opioids are preferable, maximising symptom management whilst reducing nurse and community staff time, use of resource limited protective equipment and exposure. Use of the parenteral route is also preferred for speed of onset of action and ability to rapidly titrate doses. If there is distressing breathlessness at rest, opioids should be combined with an anxiolytic sedative, e.g. midazolam (Table 1). Rapid titration of benzodiazepines may be needed. The primary role of oxygen is to correct hypoxaemia. There is a suggestion it may help breathlessness in severe hypoxaemia [14], but not when mild or absent.[15] In a comatose/unresponsive dying patient, oxygen can be titrated down with goal of discontinuation, while concurrently managing symptoms of breathlessness.[16]

Cough (70–80%)[5-9]: Potential mechanisms in viral respiratory infection and ARDS include inflammation, epithelial damage, mucus impaction and neuro-modulatory changes (heightened cough reflex sensitivity).[17] Evidence for effective management of acute cough is limited.[18, 19] Adequate hydration and regular small sips of water may help.[20] However, because of the likely concurrent need for breathlessness management, strong

opioids are likely to be more pragmatic in practice. High-level evidence is limited [21,22], but relatively low doses of long-acting oral morphine (5-10mg twice daily) are helpful in refractory chronic cough (Table 1).[21] Although these patients do not have chronic cough, opioids may be helpful in reducing cough alongside treatment for breathlessness. [22]

Delirium: Delirium is common in medical illness and almost universal in the last days–hours of life. Non-pharmacological management includes regular orientation, avoiding constipation, treating pain, maintaining oxygenation and avoiding urinary retention.[23] When ineffective and where the delirium is causing significant distress, pharmacological therapy with an antipsychotic (such as haloperidol) should be considered; in the context of severe breathlessness and at the end of life, a sedative anxiolytic such as levomepromazine or midazolam should be used (See Table 1) to allow rapid titration if needed.

Anxiety: Anxiety secondary to breathlessness, social isolation and fear is likely to be present to some degree in all patients with COVID-19. Severely ill patients may be particularly distressed, due to the looming possibility that their situation may rapidly worsen and that they may die, potentially compounded by the loss of support from their families, who are not allowed to visit, and receiving care from health professionals in personal protective equipment. Non-pharmacological methods such as relaxation therapy and breathing exercises are effective in mild anxiety but if patients are significantly anxious or have severe disease, the focus should be on pharmacological management. For severe disease and if people are actively dying, benzodiazepines are likely to be most effective.[24] Optimal relief may necessitate increasing depths of sedation rapidly in the last days–hours or of life, particularly when associated with severe breathlessness (See Table 1).

Psychological support

Families of patients with COVID-19 face a significant psychological burden that is often magnified by family members themselves being in isolation or under financial strain. Often, more than one member of the same family may be infected and in hospital. Visiting is likely to also be limited/prohibited. There may also be guilt over possibly transmitting the infection to their loved one. The quality of the dying experience and lack of preparation for the death are both predictors of complicated grief. Health professionals should communicate with families regularly and where possible facilitate communication between patients and their families utilising virtual technology.[25] Information leaflets should be provided for both patients and family members. If family members are unable to visit, leaflets should be emailed where possible. See Figure 1 and 2 for example leaflets that can be used and supplementary files for downloadable editable WORD documents.

Information needs and parallel planning

In patients with severe disease, there is a need for parallel planning- hoping for the best but preparing for the worst. Therefore, there is an urgent need for early and honest discussions at the time of hospitalisation/diagnosis of COVID-19 for those with a high risk of severe illness and death. Health professionals should focus on advance care planning with open and honest conversations as to what is important to the individual, to inform treatment escalation planning including resuscitation status, should they deteriorate.

Health professionals should acknowledge the distress of this complex and unique situation, and be compassionate, respectful and empathic. We must explain that active resuscitative care and symptom management do not need to be mutually exclusive, where resources allow. The focus of care should be iterative, adapting to worsening clinical status or patient expressed altered goals of care. Importantly, healthcare professionals need to be proactive in

the provision of information and ensure that if there is a significant possibility that the patient will die, that this is addressed with both the patient and family (Table 1).

Spiritual Care

The importance of spirituality in coping with uncertainty, severe disease and at the end of life is recognised.[26] Spiritual wellbeing offers some protection against end of life despair in those for whom death is imminent.[26] Hospital chaplains provide spiritual care that helps patients facing serious illness better cope with their symptoms and prognosis. Most hospitals around the world will have chaplains/representatives from all faiths in the chaplaincy office. The family of the patient or the dying patient may want to see speak to someone about their impending death. Chaplain interventions, whether or not religious, focused on comforting the patient and improving his or her well-being in the context of both his or her spiritual pain and critical illness should be explored with all patients and carers.[27] It is important that the spiritual care needs of those who are not represented by chaplaincy available or who are not religious, should also be addressed.

Conclusion

COVID-19 patients with severe disease/at the end of life have an equal right to care. Receiving adequate symptom control is a basic human right and we have a fundamental duty to relieve suffering and provide the best care with the available resources regardless of the chances of survival.

Acknowledgments

Thank you to all the patients and members of the public at the European Lung Foundation and Cicely Saunders Institute as well as Howard Almond (Action for Pulmonary Fibrosis) and Sarah Dix for their valuable contributions to drafting of the information leaflets.

This study/project is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Table 1 Example clinical guidelines for the management of the severe/dying COVID-19 patient

Symptom/need	Clinical indication	Recommendation
Distressing breathlessness at rest		<ul style="list-style-type: none"> Stat dose morphine 2.5mg SC/IV + midazolam 2.5mg SC/IV (reduce both to 1.25mg if eGFR <30 or elderly) <p><i>If continuous infusion is available:</i></p> <ul style="list-style-type: none"> Morphine 10mg + midazolam 10mg CSCI/ IV over 24hrs OR morphine 5mg + midazolam 5mg CSCI/IV over 24hrs (if eGFR <30 or in the elderly). In addition, prescribe morphine 2.5mg + midazolam 2.5mg SC/IV p.r.n. 4 hourly (1.25mg for both if eGFR<30 or in the elderly) <p><i>If continuous infusion is not available:</i></p> <ul style="list-style-type: none"> Morphine 2.5mg SC/IV + midazolam 2.5mg SC/IV 4 hourly (1.25mg for both if eGFR<30 or in the elderly). In addition, prescribe morphine 2.5mg SC/IV + midazolam 2.5mg SC/IV p.r.n. 4 hourly (1.25mg for both if eGFR<30 or in the elderly) <p>Monitor patients receiving opioids for undesirable effects, particularly nausea and vomiting, and constipation. Depending on individual circumstances, prescribe a regular or p.r.n anti-emetic and a regular laxative.</p>
Anxiety	Mild	<ul style="list-style-type: none"> Relaxation techniques and breathing exercises
	Moderate/severe	<ul style="list-style-type: none"> Midazolam 2.5mg SC/IV p.r.n (1.25mg for both if eGFR<30 or in the elderly) <p>If needed regularly, consider a CSCI/IV infusion (starting dose midazolam 10mg /24hr- reduce to 5mg if eGFR<30 or elderly)</p>
Cough		<p><i>If continuous infusion is available:</i></p> <ul style="list-style-type: none"> Morphine 10mg CSCI/ IV over 24hrs (5mg CSCI/IV if eGFR<30 or elderly) <p><i>If continuous infusion is not available:</i></p> <ul style="list-style-type: none"> Morphine 2.5mg SC/IV SC/IV 4 hourly (1.25mg for both if eGFR<30 or in the elderly). <p>Monitor patients receiving opioids for undesirable effects, particularly nausea and vomiting, and constipation. Depending on individual circumstances, prescribe a regular or p.r.n anti-emetic and a regular laxative.</p>
Fever		<ul style="list-style-type: none"> Paracetamol 1g QDS PO/IV (avoid NSAIDs unless end of life)
Delirium	Mild confusion	<ul style="list-style-type: none"> Orientation to time and place
	Delirium with distress	<ul style="list-style-type: none"> Orientation to time and place, if ineffective haloperidol 1-10mg SC/IV in 1-3 divided doses over 24hrs; maximum 10mg/day (halve all doses in elderly; maximum 5mg/day)
	Delirium/agitation at end of life	<ul style="list-style-type: none"> Stat dose levomepromazine 25 mg SC/IV or midazolam 5mg (reduce to levomepromazine 12.5mg and midazolam 2.5mg if eGFR<30 or elderly). <p>If needed regularly, consider CSCI/IV infusion (either levomepromazine 50mg or midazolam 15mg /24hr- reduce to levomepromazine 25mg or midazolam 7.5mg (if eGFR<20 or elderly).</p> <p>NB May need to titrate rapidly if ongoing agitation. Where on both opioid and sedative - titrate the sedative up for terminal delirium NOT the opioid</p>
Communication	Patients	<ul style="list-style-type: none"> “I understand that this is an emotional time, anyone would be scared/anxious...it is normal to be worried and scared.” “We do not think recovery from this illness is possible and [you/they] may die over the next [few hours / days / short weeks].” “I am very sorry that you cannot have your loved ones around you, but as you can see, you are here with us, you are not alone, we will stay with you.”
	Family	<ul style="list-style-type: none"> “What concerns you the most?” “It’s understandable you feel this way / This must be really hard for you / It is upsetting.” “Who is around to support you?” “Is there something we can do to help?” “We are concerned about the condition of your [relative] and think that they are ‘sick enough to die’”

Further information/resources available at: <https://www.vitaltalk.org/guides/covid-19-communication-skills/>, SIGN delirium guidance <https://www.sign.ac.uk/sign-157-delirium>

CSCI- continuous subcutaneous infusion; SC – subcutaneous; IV – intravenous; PO – per oral; BD – twice daily; GFR – glomerular filtration rate

Insert Figure 1 here

Figure 1: Example information sheet for COVID-19 patients admitted to hospital (downloadable WORD document available in supplementary files)

Insert Figure 2 here

Figure 2: Example information sheet for families/friends of COVID-19 patients admitted to hospital (downloadable WORD document available in supplementary files)

1. WHO-China Joint Mission. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19) 2020 [cited 2020 17 March 2020]; Available from: <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>
2. Rubin EJ, Baden LR, Morrissey S. Audio Interview: New Research on Possible Treatments for Covid-19. *New England Journal of Medicine* 2020: 382(12): e30.
3. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* 2020.
4. Guan W-j, Liang W-h, Zhao Y, Liang H-r, Chen Z-s, Li Y-m, Liu X-q, Chen R-c, Tang C-l, Wang T. Comorbidity and its impact on 1,590 patients with COVID-19 in China: A Nationwide Analysis. *medRxiv* 2020.
5. Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, Liu L, Shan H, Lei C-l, Hui DS. Clinical characteristics of coronavirus disease 2019 in China. *New England Journal of Medicine* 2020.
6. Yang X, Yu Y, Xu J, Shu H, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory Medicine* 2020.
7. Xu X-W, Wu X-X, Jiang X-G, Xu K-J, Ying L-J, Ma C-L, Li S-B, Wang H-Y, Zhang S, Gao H-N. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. *Bmj* 2020: 368.
8. Wang L, Gao Y-h, Zhang G-J. The clinical dynamics of 18 cases of COVID-19 outside of Wuhan, China. *European Respiratory Journal* 2020.
9. Zhang S, Li H, Huang S, You W, Sun H. High-resolution CT features of 17 cases of Corona Virus Disease 2019 in Sichuan province, China. *European Respiratory Journal* 2020.
10. Higginson IJ, Bausewein C, Reilly CC, Gao W, Gysels M, Dzingina M, McCrone P, Booth S, Jolley CJ, Moxham J. An integrated palliative and respiratory care service for patients with advanced disease and refractory breathlessness: a randomised controlled trial. *The Lancet Respiratory Medicine* 2014: 2(12): 979-987.
11. Gysels M, Reilly CC, Jolley CJ, Pannell C, Spoorendonk F, Bellas H, Madan P, Moxham J, Higginson IJ, Bausewein C. How does a new breathlessness support service affect patients? *European Respiratory Journal* 2015: 46(5): 1515-1518.
12. Ekström M, Nilsson F, Abernethy AA, Currow DC. Effects of opioids on breathlessness and exercise capacity in chronic obstructive pulmonary disease. A systematic review. *Annals of the American Thoracic Society* 2015: 12(7): 1079-1092.
13. Ekström M, Bajwah S, Bland JM, Currow DC, Hussain J, Johnson MJ. One evidence base; three stories: do opioids relieve chronic breathlessness? *Thorax* 2018: 73(1): 88-90.
14. Bruera E, de Stoutz N, Velasco-Leiva A, Schoeller T, Hanson J. Effects of oxygen on dyspnoea in hypoxaemic terminal-cancer patients. *The Lancet* 1993: 342(8862): 13-14.
15. Abernethy AP, McDonald CF, Frith PA, Clark K, Herndon II JE, Marcello J, Young IH, Bull J, Wilcock A, Booth S. Effect of palliative oxygen versus room air in relief of breathlessness in patients with refractory dyspnoea: a double-blind, randomised controlled trial. *The Lancet* 2010: 376(9743): 784-793.
16. Campbell ML, Yarandi H, Dove-Medows E. Oxygen is nonbeneficial for most patients who are near death. *Journal of pain and symptom management* 2013: 45(3): 517-523.
17. Atkinson SK, Sadofsky LR, Morice AH. How does rhinovirus cause the common cold cough? *BMJ open respiratory research* 2016: 3(1): e000118.
18. Malesker MA, Callahan-Lyon P, Ireland B, Irwin RS, Adams TM, Altman KW, Azoulay E, Barker AF, Birring SS, Blackhall F. Pharmacologic and nonpharmacologic treatment for acute cough associated with the common cold: CHEST Expert Panel Report. *Chest* 2017: 152(5): 1021-1037.

19. Hill AT, Gold PM, El Solh AA, Metlay JP, Ireland B, Irwin RS, Adams TM, Altman KW, Azoulay E, Barker AF. Adult outpatients with acute cough due to suspected pneumonia or influenza: CHEST Guideline and Expert Panel Report. *Chest* 2019; 155(1): 155-167.
20. Mitchell SAC, Garrod R, Clark L, Douiri A, Parker SM, Ellis J, Fowler SJ, Ludlow S, Hull JH, Chung KF. Physiotherapy, and speech and language therapy intervention for patients with refractory chronic cough: a multicentre randomised control trial. *Thorax* 2017; 72(2): 129-136.
21. Fuller RW, Karlsson JA, Choudry NB, Pride NB. Effect of inhaled and systemic opiates on responses to inhaled capsaicin in humans. *J Appl Physiol* 1988; 65(3): 1125-1130.
22. Morice AH, Millqvist E, Bieksiene K, Birring SS, Diczpinigaitis P, Ribas CD, Boon MH, Kantar A, Lai K, McGarvey L. ERS guidelines on the diagnosis and treatment of chronic cough in adults and children. *European Respiratory Journal* 2020; 55(1).
23. Network. SIG. Risk reduction and Management of delirium 2019 [cited March 2020]; Available from: <https://www.sign.ac.uk/sign-157-delirium>
24. Practice guideline for the treatment of patients with panic disorder. Work Group on Panic Disorder. American Psychiatric Association. *Am J Psychiatry* 1998; 155(5 Suppl): 1-34.
25. Lobb EA, Kristjanson LJ, Aoun SM, Monterosso L, Halkett GKB, Davies A. Predictors of Complicated Grief: A Systematic Review of Empirical Studies. *Death Studies* 2010; 34(8): 673-698.
26. McClain CS, Rosenfeld B, Breitbart W. Effect of spiritual well-being on end-of-life despair in terminally-ill cancer patients. *The Lancet* 2003; 361(9369): 1603-1607.
27. Berning JN, Poor AD, Buckley SM, Patel KR, Lederer DJ, Goldstein NE, Brodie D, Baldwin MR. A novel picture guide to improve spiritual care and reduce anxiety in mechanically ventilated adults in the intensive care unit. *Annals of the American Thoracic Society* 2016; 13(8): 1333-1342.

Information for patients admitted to hospital with COVID-19

COVID-19 can cause minor to serious illness.

You have been admitted to hospital with COVID-19 so that we can monitor your breathing. Your breathing may need to be supported with oxygen and sometimes with a ventilator (artificial breathing). We also want to help with any symptoms you may have.

This leaflet will explain what treatment you may receive, and what support will be available.

What treatment will I receive?

Alongside active measures to treat the disease, it is important that we reduce the distress that you may experience. This is done through treating your symptoms:

- *Breathlessness* can be improved by keeping as calm and relaxed as possible, but if your breathlessness gets worse, we will use medication to help with this. Morphine is the most common medicine used. Although usually given for pain, morphine can be used safely to relieve the feeling of breathlessness.
- *Cough* can also be relieved by morphine.
- *Anxiety* can be common; medicines used to help with this symptom include lorazepam and midazolam.
- *Restlessness* can occur if you develop a fever and this can be controlled using paracetamol.

All medicines will be given regularly and when you need them. If you become unable to swallow the medicine, it can be given as an injection either through a vein or under the skin.

In the most serious cases, COVID-19 can severely affect the lungs, stopping them from working normally. A ventilator may be used to move air into and out of the lungs to help you breathe. You may need to be on a ventilator for several days until your lungs are able to work properly again.

Can I decide how I am treated?

You should talk to the doctors about what is important to you. You may have preferences about how and when certain actions should be taken. For example, when to start ventilation or whether to restart the heart if it stops. The doctors will take your views into consideration together with your medical condition. Difficult decisions about your medical care may need to be made quickly if you become unwell so it is important that you let the medical team know what you want them to do. If you are unsure, please discuss this with a member of the medical team.

How can I communicate with those important to me?

There are strict isolation rules in place both outside and inside the hospital, which means that you may not be allowed visitors. Any visitors will have to wear personal protective equipment (otherwise known as PPE – face masks etc.). Where possible, ward staff will try to help you communicate with people important to you by telephone or video calls. Please let the ward staff know if you are happy for them to share information in this way and if there are specific people you wish to be kept informed.



This information leaflet was compiled by Dr Sabrina Bajwah (Cicely Saunders Institute, King's College London) with extensive input from co-authors and patient/carer groups to accompany the ERJ editorial "Managing the supportive care needs of those affected with COVID-19". Please cite this editorial as follows:

Bajwah S, Wilcock A, Towers R, Costantini M, Bausewein C, Simon ST, Bendstrup E, Prentice W, Johnson MJ, Currow DC, Kreuter M, Wells AU, Biring SS, Edmonds P, Higginson IJ (2020). Managing the supportive care needs of those affected by COVID-19. ERJ: doi (TO BE INSERTED)

Produced in April 2020

Can I get additional support?

We understand that this is a difficult time. You may want to talk about how you feel. Please ask a member of the ward staff to contact any of the following support services that are available to you:

List of local hospital support services:

Information for family/friends of those admitted to hospital with COVID-19

COVID-19 can cause minor to serious illness.

Your family member/friend has been admitted to hospital with COVID-19 so that we can monitor their breathing. Their breathing may need to be supported with oxygen and sometimes with a ventilator (artificial breathing). We will also help with any symptoms they may have.

This leaflet will explain what treatment they may receive, and what support will be available to them and you.

What treatment will they receive?

Alongside active measures to treat the disease, it is important that we reduce the distress that they may experience. This is done through treating symptoms:

- *Breathlessness* can be improved by keeping as calm and relaxed as possible, but if breathlessness gets worse, we will use medication to help with this. Morphine is the most common medicine used. Although usually given for pain, morphine can be used safely to relieve the feeling of breathlessness.
- *Cough* can also be relieved by morphine.
- *Anxiety* can be common; medicines used to help with this symptom include lorazepam and midazolam.
- *Restlessness* can occur if a fever develops and this can be controlled using paracetamol.

All medicines will be given regularly and when needed. Medicines can be given as an injection either through a vein or under the skin if necessary.

In the most serious cases, COVID-19 can severely affect the lungs, stopping them from working normally. A ventilator may be used to move air into and out of the lungs to help with breathing. A ventilator may be used for several days until the lungs are able to work properly again.

Making decisions

Difficult decisions about the care your family member/friend receives may need to be made rapidly by the medical teams. For example, when to start ventilation or whether to restart the heart if it stops. Please let the medical team know whether you have had these discussions with your family member/friend already and what they said. If you are unsure, then please discuss with one of the medical team. Many of the conversations with the doctors and nurses will have to take place on the telephone and we recognise that this will be difficult. Please make sure we have your correct contact details and let the ward staff know if you wish to be kept informed.

How can I communicate with them?

There are strict isolation rules in place both outside and inside the hospital, which means that you may not be allowed to visit. If visiting the hospital, you will have to wear personal protective equipment (otherwise known as PPE – face masks etc.). Where possible, ward staff will help you communicate by telephone or video calls.



This information leaflet was compiled by Dr Sabrina Bajwah (Cicely Saunders Institute, King's College London) with extensive input from co-authors and patient/carer groups to accompany the ERJ editorial "Managing the supportive care needs of those affected with COVID-19". Please cite this editorial as follows:

Bajwah S, Wilcock A, Towers R, Costantini M, Bausewein C, Simon ST, Bendstrup E, Prentice W, Johnson MJ, Currow DC, Kreuter M, Wells AU, Biring SS, Edmonds P, Higginson IJ (2020). Managing the supportive care needs of those affected by COVID-19. ERJ: doi (TO BE INSERTED)

Produced in April 2020

Can I get additional support?

We understand that this is a difficult time. You may want to talk about how you feel. Please ask a member of the ward staff to contact any of the following support services that are available to you:

List of local hospital support services: